

**PRODUCT MONOGRAPH**

**<sup>Pr</sup>CYPROTERONE**

**Cyproterone Acetate Tablets**

**House Standard**

**50 mg**

**Antiandrogen**

**AA PHARMA INC.  
1165 Creditstone Road Unite 1  
Vaughan, Ontario  
L4K 4N7**

**DATE OF REVISION:  
November 25, 2014**

**Control No: 163127**

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS .....	3
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS.....	8
DRUG INTERACTIONS .....	9
DOSAGE AND ADMINISTRATION.....	10
OVERDOSAGE.....	11
ACTION AND CLINICAL PHARMACOLOGY.....	11
STORAGE AND STABILITY .....	12
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	12
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>13</b>
PHARMACEUTICAL INFORMATION .....	13
CLINICAL TRIALS .....	13
DETAILED PHARMACOLOGY .....	19
TOXICOLOGY.....	21
REFERENCES.....	26
<b>PART III: CONSUMER INFORMATION .....</b>	<b>29</b>

**CYPROTERONE**  
**Cyproterone Acetate Tablets**

**House Standard**

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form, Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients</b>
oral	Tablets, 50 mg	<i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

**INDICATIONS AND CLINICAL USE**

CYPROTERONE (cyproterone acetate) is indicated for:

- the palliative treatment of patients with advanced prostatic carcinoma

**CONTRAINDICATIONS**

- Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section.
- Liver disease and hepatic dysfunction
- Dubin Johnson syndrome, Rotor syndrome
- Previous or existing liver tumors (only if these are not due to metastases from carcinoma of the prostate)
- Presence or history of meningioma
- Wasting diseases (with the exception of inoperable carcinoma of the prostate)
- Severe chronic depression
- Existing thromboembolic processes

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

CYPROTERONE should be prescribed and managed by a qualified physician experienced in the use of hormonal therapy in prostate cancer. The following are clinically significant adverse events:

- Hepatotoxicity with acute hepatic failure (see **Hepatic/Biliary/Pancreas, Hepatotoxicity**).

### General

**Concomitant Alcohol:** Alcohol may reduce the antiandrogenic effect of CYPROTERONE in hypersexuality. The relevance of this in prostatic carcinoma is not known; however, it would be prudent to inform the patients that the use of alcohol during CYPROTERONE therapy is not advisable.

**Physical Performance:** Patients should be informed that fatigue and lassitude are common in the first few weeks of therapy, but usually become much less pronounced from the third month on. Marked lassitude and asthenia necessitate special care when driving or operating machinery.

**Concomitant Use With a GnRH Agonist or Orchiectomy:** Based on a retrospective meta-analysis, long-term combination therapy of CYPROTERONE with either orchiectomy or a GnRH agonist as treatment of patients with advanced prostate cancer may result in a 5-year survival disadvantage compared to castration alone.

### Carcinogenesis and Mutagenesis

Cyproterone acetate showed a potential to initiate and/or promote liver tumor formation in rodents. Very rare cases of benign and malignant liver tumors have been observed in patients receiving cyproterone acetate.

**Meningioma:** The occurrence of meningiomas (single and multiple) has been reported in association with long-term use (years) of cyproterone acetate at doses of 25 mg/day and above. In a retrospective cohort study using data from a primary care database, meningiomas were reported very rarely in patients treated with cyproterone acetate for prostate cancer after several months of treatment; in these cases, causality was not established. If a patient treated with CYPROTERONE is diagnosed with a meningioma, treatment with CYPROTERONE must be stopped. Patients with prehistory or presence of meningioma should not be treated with CYPROTERONE (see **CONTRAINDICATIONS**).

**Antiandrogen Withdrawal Syndrome:** In some patients with metastatic prostate cancer,

antiandrogens (steroidal or nonsteroidal) may promote, rather than inhibit, the growth of prostate cancer. A decrease in PSA and/or clinical improvement following the discontinuation of antiandrogens has been reported. It is recommended that patients prescribed an antiandrogen, who have PSA progression, should have the antiandrogen discontinued immediately and be monitored for 6 to 8 weeks for a withdrawal response prior to any decision to proceed with other prostate cancer therapy.

**Gynecomastia:** Benign nodules (hyperplasia) of the breast have been reported; these generally subside 1 to 3 months after discontinuation of therapy and/or after a reduction of dosage. The reduction of dosage should be weighed against the risk of inadequate tumor control.

### **Endocrine and Metabolism**

**Adrenocortical Function:** Suppression of adrenocortical function tests have occurred in patients receiving cyproterone acetate and preclinical data also revealed a suppression of adrenal gland due to the administration of cyproterone acetate (see **TOXICOLOGY**).

Reduced response to endogenous ACTH was noted by metyrapone test; furthermore, reduced ACTH and cortisol blood levels determined by the Mattingly method were also found. It is therefore recommended that adrenocortical function tests should be monitored periodically by serum cortisol assay.

**Diabetes:** CYPROTERONE may impair carbohydrate metabolism. Parameters of carbohydrate metabolism, fasting blood glucose, and glucose tolerance tests, should be examined carefully in all patients and particularly in all diabetics before and regularly during therapy with CYPROTERONE.

Strict medical supervision is necessary if the patient suffers from diabetes, because the requirement for oral antidiabetics or insulin can change during CYPROTERONE treatment.

**Metabolic Effects:** Fluid retention, hypercalcemia and changes in plasma lipid profile may occur. Accordingly, CYPROTERONE should be used with caution in patients with cardiac disease.

**Nitrogen Balance:** A negative nitrogen balance is usual at the start of therapy, but does generally correct itself within 3 months of continued therapy.

### **Hematologic**

**Hematology:** Hypochromic anemia has been observed rarely during therapy with cyproterone acetate. Regular hematological assessment is recommended.

**Thromboembolism:** Clinical investigations have shown that when cyproterone acetate is used alone it has a minor effect on blood clotting factors. However, when cyproterone acetate was combined with ethinyl estradiol, changes were found in increased coagulation capability.

The occurrence of thromboembolic events has been reported in patients using,

cyproterone acetate although a causal relationship has not been established. Patients with previous arterial or venous thrombotic / thromboembolic events (eg, deep venous thrombosis, pulmonary embolism, myocardial infarction) or with a history of cerebrovascular accidents or with advanced malignancies are at increased risk of further thromboembolic events. CYPROTERONE should be discontinued at the first sign of thrombophlebitis or thromboembolism, and the patient should be carefully re-evaluated if manifestations of thrombotic disorder occur: thrombophlebitis, cerebrovascular complications, retinal thrombosis, or pulmonary embolism.

In patients with inoperable carcinoma of the prostate, presenting with a history of thromboembolic processes or suffering from sickle cell anemia or from severe diabetes with vascular changes, a careful risk:benefit evaluation must be carried out in each individual case before CYPROTERONE is prescribed.

### **Hepatic/Biliary/Pancreas**

**Hepatotoxicity:** Direct hepatic toxicity, including jaundice, hepatitis, and acute hepatic failure has been observed in patients treated with cyproterone acetate. At daily doses of 100 mg and above cases with fatal outcome have also been reported. Most reported fatal cases were in men treated with cyproterone acetate for prostatic cancer. Hepatotoxicity is dose-related and develops, usually, a few weeks to several months after cyproterone treatment has begun. Liver function tests should be performed pretreatment, at regular intervals during treatment, and whenever any symptoms or signs suggestive of hepatotoxicity occur. If hepatotoxicity is confirmed, CYPROTERONE should be withdrawn. Benefit and risk should be evaluated carefully if any drug(s) with known hepatotoxicity is to be used concurrently with CYPROTERONE. CYPROTERONE should not be used in patients with prior history or existing hepatic disease (see **CONTRAINDICATIONS**).

In very rare cases, benign and malignant liver tumors which may lead to life-threatening intra-abdominal hemorrhage have been observed after the use of cyproterone acetate. If severe upper abdominal complaints, liver enlargement, or signs of intra-abdominal hemorrhage occur, a liver tumor should be included in the differential-diagnostic considerations.

### **Psychiatric**

**Depression:** Cyproterone acetate therapy has occasionally been associated with an increased incidence of depressive mood changes, especially during the first 6 to 8 weeks of therapy. Similar mood changes have also been seen following surgical castration and are considered to be due to androgen deprivation. Patients with tendencies to depressive reaction should be carefully observed.

### **Respiratory**

**Shortness of Breath:** A sensation of shortness of breath was commonly reported in patients treated with 300 mg/day cyproterone acetate. Patients with pre-existing pulmonary dysfunction are most likely to be affected.

## **Sexual Function/Reproduction**

**Inhibition of Spermatogenesis:** The sperm count and the volume of ejaculate are reduced at oral doses of 50 to 300 mg per day. Infertility is usual, and there may be azoospermia after 8 weeks of therapy, which is associated with atrophy of seminiferous tubules.

Follow-up examinations on discontinuation of therapy have shown these changes to be reversible.

Spermatogenesis usually reverts to its previous level about 3 to 5 months after stopping CYPROTTERONE, or in some patients, after up to 20 months. Production of abnormal spermatozoa during cyproterone acetate therapy has been observed; their relationship to abnormal fertilization or malformed embryos is not known.

## **Skin**

CYPROTTERONE therapy may cause a reduction of sebum production leading to dryness of the skin and transient patchy loss of body hair.

## **Special Populations**

### ***Pregnant Women***

Treatment with CYPROTTERONE is not indicated for use in women.

### ***Nursing Women***

Treatment with CYPROTTERONE is not indicated for use in women.

### ***Pediatrics***

CYPROTTERONE is not recommended for use in children and adolescents below 18 years of age. CYPROTTERONE must not be given before the conclusion of puberty since an unfavorable influence on longitudinal growth and the still unstabilized axes of endocrine function cannot be ruled out.

## **Monitoring and Laboratory Tests**

It is recommended that patients prescribed an antiandrogen, who have PSA progression, should have the antiandrogen discontinued immediately and be monitored for 6 to 8 weeks for a withdrawal response prior to any decision to proceed with other prostate cancer therapy.

With the potential for adrenal gland suppression, it is recommended that adrenocortical function tests should be monitored periodically by serum cortisol assay.

Parameters of carbohydrate metabolism, fasting blood glucose, and glucose tolerance tests, should be examined carefully in all patients and particularly in all diabetics before and regularly

during therapy with CYPROTERONE.

During treatment with CYPROTERONE, serum electrolytes and complete blood counts should be performed regularly. Liver function tests should be performed pretreatment, at regular intervals during treatment, and whenever any symptoms or signs suggestive of hepatotoxicity occur.

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

The adverse events associated most frequently with the use of cyproterone acetate are those related to the hormonal effects of the drug. These reactions usually disappear upon discontinuation of therapy or reduction of dose: decreased libido, breast enlargement, breast tenderness, benign nodular hyperplasia of the breast, galactorrhea, gynecomastia, abnormal spermatozoa, impotence, and inhibition of spermatogenesis.

The most serious adverse drug reactions (ADRs) in patients receiving cyproterone acetate are hepatic toxicity, benign and malignant liver tumors which may lead to intra-abdominal hemorrhage, and thromboembolic events.

As with other antiandrogenic treatments, long-term androgen deprivation with cyproterone acetate may lead to osteoporosis.

Other adverse events which have been reported are listed below:

**Cardiovascular System:** hypotension, tachycardia, heart failure, syncope, myocardial infarct, hemorrhage, cerebrovascular accident, cardiovascular disorder, retinal vascular disorder, embolus, pulmonary embolism, superficial and deep thrombophlebitis, thrombosis, retinal vein thrombosis, phlebitis, vascular headache, shock, pulmonary oil microembolism, vasovagal reactions.

**Gastrointestinal System:** constipation, diarrhea, indigestion, anorexia, nausea, vomiting, cholestatic jaundice, cirrhosis of liver, hepatic coma, hepatitis, hepatoma, hepatomegaly, jaundice, liver carcinoma, liver failure (for further information see **WARNINGS AND PRECAUTIONS - Hepatic/Biliary/Pancreas**), abnormal liver function test, liver necrosis, pancreatitis, glossitis.

**Hematology:** increased fibrinogen, decreased prothrombin, thrombocytopenia, anemia (for further information see **WARNINGS AND PRECAUTIONS - Hematologic**), hemolytic anemia, hypochromic anemia, normocytic anemia, leukopenia, leukocytosis.

**Metabolism:** negative nitrogen balance, decreased response to ACTH, hyperglycemia, lowered cortisol, hypercalcemia, increased SGOT, increased SGPT, increased creatinine, hypernatremia, edema, weight gain, weight loss, diabetes mellitus.

**Musculoskeletal System:** myasthenia, osteoporosis.

**Central Nervous System:** fatigue, lassitude, weakness, hot flashes, increased sweating, aphasia, coma, depression, dizziness, encephalopathy, hemiplegia, personality disorder, psychotic depression, abnormal gait, headache, temporary restlessness.

Meningiomas (single and multiple) have been reported in association with long-term use (several years) of cyproterone acetate. In a retrospective cohort study using data from a primary care database, meningiomas were reported very rarely in patients treated with cyproterone acetate for prostate cancer after several months of treatment; in these cases, causality was not established (see **WARNINGS AND PRECAUTIONS - Carcinogenesis and Mutagenesis**).

**Respiratory System:** asthma, increased cough, dyspnea, hyperventilation, respiratory disorder, shortness of breath on effort (see **WARNINGS AND PRECAUTIONS - Respiratory**), lung fibrosis.

**Skin:** eczema, urticaria, erythema nodosum, exfoliative dermatitis, rash, maculopapular rash, dryness of the skin, pruritus, alopecia, hirsutism, skin discoloration, photosensitivity reactions, scleroderma.

**Sensory System:** ear disorder, optic atrophy, optic neuritis, abnormality of accommodation, abnormal vision, blindness, retinal disorder.

**Urogenital System:** enlarged uterine fibroids, uterine hemorrhage, increased urinary frequency, bladder carcinoma, kidney failure, hematuria, urate crystalluria, urine abnormality.

**Other:** ascites, allergic reaction, asthenia, chills, fetal chromosome abnormality, death, fever, hernia, malaise, injection site reaction.

Adverse reactions are rarely of sufficient severity to require dosage reduction or discontinuation of treatment.

If reactions are severe, it may be beneficial to reduce the dosage.

## **DRUG INTERACTIONS**

### **Overview**

Although clinical interaction studies have not been performed, since this drug is metabolized by CYP3A4, it is expected that ketoconazole, itraconazole, clotrimazole, ritonavir, and other strong inhibitors of CYP3A4 inhibit the metabolism of cyproterone acetate. On the other hand, inducers of CYP3A4 such as rifampicin, phenytoin, and products containing St. John's wort may reduce the levels of cyproterone acetate.

Based on in vitro inhibition studies, an inhibition of the cytochrome P450 enzymes CYP2C8, 2C9, 2C19, 3A4, and 2D6 is possible at high therapeutic cyproterone acetate doses of 300 mg daily. In addition, cyproterone acetate was also shown to increase the enzymatic activity of CYP1A2 and CYP2E1 in vitro. Caution should be exercised when CYPROTERONE is to be co-administered with a

substrate of the P450 enzymes.

The risk of statin-associated myopathy or rhabdomyolysis may be increased when those HMGCoA inhibitors (statins) which are primarily metabolized by CYP 3A4 are coadministered with high therapeutic cyproterone acetate doses, since they share the same metabolic pathway.

### **Drug-Drug Interactions**

Interactions with other drugs have not been established.

### **Drug-Food Interactions**

Interactions with food have not been established.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

#### ***Patients with Hepatic Impairment***

The use of CYPROTERONE is contraindicated in patients with liver diseases and/or with hepatic dysfunction.

#### ***Patients with Renal Impairment***

A pharmacokinetic study in patients with renal impairment has not been conducted. As 33% of cyproterone acetate is excreted via the kidney, caution should be taken when CYPROTERONE is administered in this patient population.

### **Recommended Dose and Dosage Adjustment**

**Oral Tablets:** The usual daily initial and maintenance dose of CYPROTERONE (cyproterone acetate) is 4 to 6 tablets (200-300 mg) divided into 2 to 3 doses and taken with some liquid after meals.

The maximum daily dose is 300 mg.

After orchiectomy, a lower daily dose of 2 to 4 tablets (100-200 mg) is recommended.

CYPROTERONE therapy should not be interrupted nor the dosage reduced after remission or improvement occurs.

Because of their pharmacokinetic properties, cyproterone acetate (oral) and cyproterone acetate (injectable) can be interchanged in the course of long-term treatment. The dosage may be reduced if side effects are intolerable, but should be kept within the oral range of 2 to 6 tablets daily (100-300 mg).

## **OVERDOSAGE**

For management of a suspected drug overdose, contact your regional Poison Control Centre.

There have been no reports of fatal overdosage in man with cyproterone acetate. There are no specific antidotes and treatment should be symptomatic. If oral overdosage is discovered within two to three hours, gastric lavage can safely be used if indicated.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

CYPROTERONE (cyproterone acetate) is a steroid which clinically demonstrates two distinct properties:

- a) Antiandrogenic: Cyproterone acetate blocks the binding of dihydrotestosterone - the active metabolite of testosterone - to the specific receptors in the prostatic carcinoma cell.
- b) Progestogenic/antigonadotrophic: Cyproterone acetate exerts a negative feed-back on the hypothalamo-pituitary axis, by inhibiting the secretion of LH leading to diminished production of testicular testosterone.

### **Pharmacokinetics**

#### ***Absorption***

The absorption of cyproterone acetate following oral administration is complete. Peak plasma levels are reached 3 to 4 hours after administration. Plasma levels fall rapidly during the first 24 hours as a result of tissue distribution and excretion, and plasma half-life was  $38 \pm 5$  hours.

#### ***Metabolism***

The principal metabolite identified was 15 -hydroxy-cyproterone acetate.

#### ***Excretion***

Most of the cyproterone acetate is excreted unchanged in the feces (60%) or urine (33%) within 72 hours.

Cyproterone acetate is eliminated with the urine mainly in the form of unconjugated metabolites and with the bile (feces) in the form of glucuronidized metabolites.

### **STORAGE AND STABILITY**

CYPROTERONE should be stored store at room temperature (15-30°C).

### **DOSAGE FORMS, COMPOSITION AND PACKAGING**

CYPROTERONE Cyproterone Tablets BP contain 50 mg of cyproterone acetate.

CYPROTERONE Tablets also contain the following non-medicinal ingredients: colloidal silicon dioxide, croscarmellose sodium and magnesium stearate.

CYPROTERONE (cyproterone acetate) 50 mg tablets BP are off-white, round, flat-faced tablets with bevelled-edges, embossed "CYP" over "50" scored on one side.

CYPROTERONE Tablets are available in bottles of 100, and unit dose packages of 60 and 100 tablets.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

**Proper Name:** Cyproterone acetate

**Chemical Names:**

- 1) 3'*H*-Cyclopropa[1,2]pregna-1,4,6-triene-3,20-dione, 17-(acetyloxy)-6-chloro-1,2-dihydro-, (1,2)-;
- 2) 6-Chloro-1,2-dihydro-17-hydroxy-3'*H*-cyclopropa[1,2]-pregna-1,4,6-triene-3,20-dione acetate.

#### Structural Formula:



**Molecular Formula:** C<sub>24</sub>H<sub>29</sub>ClO<sub>4</sub>

**Molecular Weight:** 416.94

#### Description:

White or almost white, odourless, crystalline powder. Insoluble in water, very soluble in chloroform, freely soluble in acetone and soluble in methanol. Melting point is about 210°C.

### CLINICAL TRIALS

#### Comparative Bioavailability

A standard, randomized, two-way crossover study was conducted in 18 healthy, adult, male volunteers to evaluate the relative bioavailability of single oral doses of CYPROTERONE 50 mg

Tablets and Androcur® 50 mg Tablets. The mean pharmacokinetic parameters of these subjects are summarized Table 1.

**Table 1:**

<b>Summary Table of the Comparative Bioavailability Data Cyproterone Acetate (Dose: 1 x 50 mg) From Measured Data</b>				
	<b>Geometric Mean Arithmetic Mean (CV%)</b>			
<b>Parameter</b>	<b>CYPROTERONE 50 mg Tablets</b>	<b>ANDROCUR®† 50 mg Tablets</b>	<b>Ratio of Geometric Means (%)**</b>	<b>90% Confidence Interval</b>
AUC <sub>0-72</sub> (ng·h/mL)	1966.3023 2010.819 (21)	2108.7500 2148.491 (18)	93.2%	88.8% - 97.9%
AUC <sub>I</sub> (ng·h/mL)	2566.5294 2662.606 (27)	2724.9591 2806.800 (24)	94.2%	88.6% - 100.1%
C <sub>MAX</sub> (ng/mL)	103.1616 106.854 (31)	128.6780 134.074 (28)	80.2%	70.6% - 91.1%
T <sub>MAX</sub> * (h)	3.02 (61)	2.50 (32)	--	
T <sub>1/2</sub> * (h)	37.60 (31)	38.21 (26)	--	

\* Arithmetic means (CV%); \*\* Based on the least squares estimate;

† Androcur® is manufactured by Berlex Canada Inc. and was purchased in Canada.

A total of 24 studies have been conducted with cyproterone acetate in patients requiring palliative treatment for advanced prostatic carcinoma. Worldwide, more than 1,000 patients have participated in these studies, which included several large multicentre trials in addition to the important comparative multicentre trial conducted by the European Cancer Oncology Group. North American experience has been accumulated in the U.S. by Drs. Scott (Johns Hopkins Hospital, Baltimore), Geller (Mercy Hospital & Medical Center, San Diego), and by Drs. Wein and Murphy (Hospital of the University of Pennsylvania, Philadelphia).

## **Study Demographics and Trial Design**

### ***Patients and Stage of Disease***

As shown in Table 2, more than 90% of the patients treated with cyproterone acetate had stage C advanced prostatic carcinoma, or stage D1 or D2 prostatic carcinoma with metastasis.

**Table 2: Patients**

Stage	No. of Patients
A or B	18
C	174
C or D	502
D	349
Not specified	39
Total	1082

The majority of patients (75%) had had no therapy prior to treatment with cyproterone acetate. A large group of patients had received various types of estrogen therapy, but had proven to be refractory or unable to tolerate the drug. A few patients had undergone an orchiectomy or had received radiation therapy (Table 3).

**Table 3: Previous Therapy**

Previous Therapy	No. of Patients
None	809
Orchiectomy	76
Estrogen	253
Radiation	16

### *Dosage and Administration*

The oral route of administration of cyproterone acetate was employed in 910 patients (84%) As shown in the table below (Table 4), the daily oral dose varied considerably from study to study and from patient to patient. However, most patients were treated with doses ranging from 200 to 300 mg/day. In orchiectomized patients, the daily dose was generally reduced by about 50% to a range of 100 to 200 mg/day orally.

**Table 4: Dose of cyproterone acetate**

Entity	Route	Dose	No. of Patients
cyproterone acetate	Oral	100 mg/day	15
		200 mg/day	197
		250 mg/day	135
		300 mg/day	114
		100-300 mg/day	449

Only 32 patients (3%) received concomitant drug therapy with cyproterone acetate. No other patients received concomitant drugs, but 521 patients (48%) underwent an orchiectomy (Table 5).

**Table 5: Concomitant Therapy**

Concomitant Therapy	No. of Patients
None	529
Estrogen (DES 0.1 mg)	32
Orchiectomy	521

## Study Results

### Effect on Serum Testosterone and Prostatic Acid Phosphatase (PAP)

**Table 6: Effect on Serum Testosterone and Prostatic Acid Phosphatase (PAP)**

Parameter	No. of Studies	Result
Serum testosterone	7	70-90% reduction
Prostatic acid phosphatase	11	Normalization in 90% of responding patients

The effect of cyproterone acetate on serum testosterone was monitored in 7 studies (Table 6). Serum testosterone was rapidly reduced following daily oral doses of 200 to 300 mg, with castrate levels being achieved within 1 to 4 weeks. The reduction was usually in the order of 70% to 90%; the greatest percent reduction occurred when cyproterone acetate was combined with estrogen.

Results of PAP evaluations consistently showed a normalization of values within a very short time in responding patients. Similarly, when there were signs of progressing metastasis, PAP values again deviated from normal levels.

### *Effect on Primary Tumor*

The effect of cyproterone acetate on the primary tumor was assessed in a total of 678 patients. Of these, 489 were previously untreated; the primary tumor was reduced in 318 of these (65%) and was stabilized in another 69 (14%). Thus, the overall positive response rate in this group was 79% (Table 7).

A significant, though smaller, percentage (59%) of estrogen-refractory patients also exhibited a positive result.

**Table 7: Effect on Primary Tumor**

Patient Group	Number	Response of Primary Tumor		Total With Positive Effect
		Reduced	Stabilized	
Previously untreated	489			
Estrogen refractory	189	318 (65%)	69 (14%)	387 (79%)
		112 (59%)	-	112 (59%)

### *Effect on Metastasis*

As shown in Table 8, metastasis was reduced in 31% of 216 evaluable patients who had not previously been treated, but in only 13% of the evaluable estrogen-refractory patients. The progression of metastases appeared to be time-dependent. Despite reduced serum testosterone

levels, metastases progressed over a period of several months to years, even in patients who were initially stabilized. The major cause of death during therapy with cyproterone acetate was the progression of metastases and not the primary tumors.

**Table 8: Effect on Metastases**

Patient Group	Number	Response of Metastases		Total with Positive Effect
		Reduced	Stabilized	
Previously untreated	216			
Estrogen refractory	71	67 (31%)	82(39%)	149(70%)
		10 (13%)	7(10%)	17(23%)

***Effect on Pain***

Table 9 illustrates the incidence of pain relief reported in each of 13 studies. Pain relief was noted in approximately 50% to 80% of patients receiving treatment with cyproterone acetate. The effect of cyproterone acetate on pain generally paralleled its effect on metastases. As long as metastases remained improved or stabilized, the analgesic requirement was also reduced. Renewed analgesic requirements were frequently indicative of metastatic progression.

**Table 9: Pain Relief**

Investigator	Incidence of Pain Relief
Dr. Bracci	172/216
Dr. Giuliani	12/16
Dr. Smith	12/25
Dr. Scott	8/10
Dr. Geller	8/10
Dr. Mauermayer	38/58
Dr. Wein	13/24
Dr. Tveter	2/6
Dr. Di Silverio	13/20
Dr. Ah-Lan	9/16
Dr. Pescatore	12/16
Dr. Hermabessiere	2/4
Dr. Bruchovsky	15/24
<b>Total</b>	<b>316/425 = 74%</b>

***Subjective and Objective Responses***

A general improvement in the subjective assessment of the quality of life was achieved in 70% of the 367 evaluable patients (Table 10).

The objective evaluations of remissions shown in Table 10 were based on ECOG criteria. The best results were obtained when cyproterone acetate was used in combination with orchiectomy. One study revealed that more than 1/3 of the patients treated with cyproterone acetate achieved a complete or partial remission for 3 to 5 years. The Canadian study found that a complete or partial remission was still evident in 75% of the patients after one year of treatment.

**Table 10: Subjective and Objective Responses**

<b>Subjective Responses</b>			
<b>No. Evaluable Patients</b>		<b>No. Improved<sup>a</sup></b>	
367		255(70%)	
<b>Objective Responses (ECOG Criteria)</b>			
<b>Treatment</b>	<b>Patient Group</b>	<b>No. of Patients</b>	<b>No. With Complete or Partial Remissions</b>
cyproterone acetate	Previously untreated	270	134 (50%)
cyproterone acetate	Estrogen-refractory	77	31 (44%)
cyproterone acetate Orchiectomy	Previously untreated and/or estrogen- refractory	274	154 (60%)

a Based on criteria of general improvement in quality of life (ie, weight gain, pain relief, etc.)

### ***Survival Rate***

**Table 11: Survival Rate**

<b>Investigator</b>	<b>No. of Patients</b>	<b>Stage</b>	<b>Duration of Treatment</b>	<b>Survival</b>	
				<b>cyproterone acetate</b>	<b>Estrogen</b>
Dr. Mauermayer	58	C or D	2-5 years	38/58 (70%)	-
Dr. Wein	55	A (7) C (25) D (23)	4 years	39/55(70%)	-
Dr. Bracci	216	C or D	5 years	138/216 (64%)	-
Dr. Di Silverio	20	D	Up to 38 months	3/20 (15%)	-
Dr. Giuliani	68	C	5 years	30/68 (44%)	31%
Dr. Giuliani	38	D	3 years	10/38 (27%)	10%
Dr. Jacobi	51	C or D	2 years	18/40 (45%)	-
Dr. Pavone	103	C or D	3.5 ó 5 years	42/103 (41%)	41%
Dr. Bruchovsky	29	D	9 ó 15 months	23/29 (80%)	-

As shown in Table 11 above, 5-year survival rates ranged from 41% to 64%. The 3-year rate for stage D patients was 27% and 1- to 2-year rates varied from a low of 15% up to a high of 80%. These survival rates generally represented an improvement over results previously obtained with estrogen therapy.

## DETAILED PHARMACOLOGY

### Animal Pharmacology

#### *Antiandrogenic Effects*

Cyproterone acetate at doses of 10 or 50 mg/kg inhibits the effects of endogenously produced and exogenously administered androgens at the prostate by means of competitive inhibition.

In mice and dogs, cyproterone acetate induces a dose-dependent atrophy of the accessory sex glands, the prostate, seminal vesicles, and preputial glands.

Spermatogenesis is inhibited in a dose-related manner; however, the atrophy in the Leydig cells are slight.

In the rat the start of puberty is prevented or delayed. Cyproterone acetate inhibits the physiological closure of the epiphyseal cartilages and bone maturation.

It impairs the function of the sebaceous glands, and the thickness of the epidermis decreases.

The treatment of pregnant animals with cyproterone acetate leads to developmental disturbances in male fetuses. Testosterone-dependent differentiation processes are affected: signs of feminization of varying degrees of severity develop.

#### *Progestogenic and Antigonadotrophic Effect*

On subcutaneous injections a total dose of 0.003 mg cyproterone acetate is about 100 times stronger than progesterone in the maintenance of pregnancy (Clausberg test). Like all potent progestogens, cyproterone acetate has antigonadotrophic properties which can be demonstrated in the parabiosis test, the testicular inhibition test in infantile rats, and by the inhibition of ovulation.

#### *Pharmacokinetic Studies in Animals*

Pharmacokinetic studies have been carried out in a number of animal species (rats, rabbits, dogs, and monkeys) using either methylene-<sup>14</sup>C- or carboxy-<sup>14</sup>C-labelled cyproterone acetate.

Cyproterone acetate is absorbed at most dose levels tested except in high doses. Peak plasma levels are usually obtained within 1 to 4 hours of oral dosing. Because of its lipophilic character, cyproterone acetate is taken up and concentrated in the liver and fatty tissues in all animal species. Cyproterone acetate is not hydrolysed, and mainly cyproterone acetate and the metabolite 15 -hydroxycyproterone acetate are found in the tissues and in plasma. The elimination half-life of cyproterone acetate is slow in most species (1-2 days), in a ratio of 4:6 with urine and feces; an exception is the dog, which excretes cyproterone acetate in 1 to 3 days. On repeated daily dosing, cyproterone acetate shows limited rise, and plasma levels can be taken as a reliable index of the concentrations of cyproterone acetate in the body. Cyproterone acetate passes the placental barrier, but only reaches the fetus in low concentrations. The

pharmacokinetics, biotransformation, and metabolic spectra of cyproterone acetate are similar in man and the rhesus monkey.

## **Human Pharmacology**

### ***Antiandrogenic Effect***

The following actions which are associated with the antiandrogenic effects have been described in man: reduction of sexual drive; inhibition of spermatogenesis; palliative effect in prostatic carcinoma; inhibition of sebaceous gland activity; suppression of signs of androgenization in women; inhibition of premature genital development in children; and other associated symptoms.

### ***Progestogenic and Antigonadotrophic Effect***

Cyproterone acetate in man is also a potent progestogen and has an antigonadotrophic effect. It intervenes with the hypothalamo-pituitary pathway, causing an inhibition of increased secretion of LH, and a decrease in gonadal testicular androgens.

Thus, unlike pure antiandrogens, cyproterone acetate does not cause a compensatory increase in androgen secretion.

### ***Other Endocrine Effects***


No distinct influence on the 17-ketosteroids, 17-ketogenic steroids or on total estrogens in the 24-hour urine has been observed in male patients. On fluorometric determination of urinary cortisol, the value apparently increases because the cyproterone acetate eliminated with the urine is also measured. Simultaneously, cyproterone acetate also reduces the reaction of the adrenal cortex to exogenous ACTH in patients; the baseline cortisol and ACTH values may also be reduced.

### ***Pharmacokinetics***

A bioavailability study was performed in 5 male volunteer subjects receiving a single oral dose of 50 mg <sup>14</sup>C-cyproterone acetate tablets.


Results of the study showed that cyproterone acetate is absorbed slowly, but completely (100%), from the gastrointestinal tract. The maximum plasma level was reached 3 to 4 hours after ingestion. The mean plasma levels were 700 nmol/L (=290µg/L) cyproterone acetate or, including the radioactivity of metabolites, 960 nmol/L (=400µg/L) cyproterone acetate equivalent.

The plasma levels fell quickly up to 24 hours after administration because of extensive tissue distribution. The half-life of cyproterone acetate in plasma was calculated as  $38 \pm 5$  hours (see Figure 1).



**Figure 1: Relationship of Unchanged Cyproterone Acetate to the Total  $^{14}\text{C}$ -labelled Substance (Cyproterone Acetate + Metabolites) in the Plasma of a Male Subject Following Oral Administration of 50 mg  $^{14}\text{C}$ -cyproterone Acetate**

On oral administration cyproterone acetate was eliminated with a half-life of  $38 \pm 2$  hours. After 10 days,  $33 \pm 6\%$  of the dose could be recovered in the urine and  $60 \pm 8\%$  in the feces (see Figure 2).



**Figure 2: Elimination (% of Dose) Following Oral Administration of 50 mg  $^{14}\text{C}$ -cyproterone Acetate in Male Subjects. Mean Values  $\pm$  Standard Deviation (n=5)**

## TOXICOLOGY

Cyproterone acetate has been found at low doses of 2 to 10 mg/kg to cause liver abnormalities in dogs and rats in the form of proliferative liver changes including increased liver weight, liver cell hypertrophy with an increase in the smooth endoplasmic reticulum, and a rise in the serum glutamic pyruvic transaminase (SGPT). At high doses of 50 to 100 mg/kg, nodular hepatic hyperplasia and hepatomas have also been observed.

In repeat-dose studies conducted in rats (12 weeks) and dogs (54 weeks) with oral administration of cyproterone acetate, decreased adrenal weights in rats at 4.0 mg/kg/day and dogs at 10 mg/kg were noted. Marked atrophy of zona fasciculata and of zona reticularis with preservation of zona glomerulosa was also observed in the adrenal glands of all treated dogs.

## Acute Toxicity

The LD<sub>50</sub> after single application of cyproterone acetate was as follows:

**Table 12: LD<sub>50</sub> After Single Application of Cyroterone Acetate**

Animal species	Oral (mg/kg)	Subcutaneous (mg/kg)	Intraperitoneal (mg/kg)	Intramuscular (mg/kg)
Mouse	>6000	>5000	>4000	-
Rat	>4000	1500	1000	-
Dog	>3000	-	-	>100 (approx.)

On the basis of the above LD<sub>50</sub> values, cyproterone acetate can be considered practically nontoxic following single dose administration. The maximum intramuscular doses were also tolerated without symptoms in the dog, with exception of local tolerance manifestation.

## Repeated Dose Toxicity

Repeat-dose toxicity studies revealed pathological changes in the liver, reproductive organs, adrenal glands, abnormal laboratory tests, and neoplasms of various tissues and organs in the animal species tested.

**Table 13: Chronic Toxicity Studies**

Animal Species	Dosage and Duration	Mortality and Clinical and Laboratory Observations	Necropsy and Histopathology
Rats 35/sex/dose	0; 10; 50, and 250 mg/kg 78 weeks oral	250 mg/kg: marked increase in mortality rate. 50 and 250 mg/kg: 40-50% decrease in body weight gain. SGPT increase: males 10 and 250 mg/kg; females 50 mg/kg. BUN increase: males 50 and 250 mg/kg. Cholesterol increase: all treated groups.	Dose-related increase in liver weights. Increase thyroid weight except for low dose males. Dose-related decrease in gonads, adrenal, prostate, seminal vesicle, and uterus weights. Histopathology: toxic manifestation in liver and kidneys - less at 10 mg/kg, more extensive at 50 and 250 mg/kg. Changes included: yellow nodules and mottling of liver (including liver cell hyperplasia and liver cell adenomas and endoplasmic inclusion bodies), discolored kidneys with rough surfaces.
Rats	0; 0.04; 0.4,	No drug-related mortality.	2 mg/kg increased

<b>Animal Species</b>	<b>Dosage and Duration</b>	<b>Mortality and Clinical and Laboratory Observations</b>	<b>Necropsy and Histopathology</b>
60/sex/dose	and 2 mg/kg 104 weeks oral	Dose-related decrease in body weight gains in males and increase in females. Food consumption reduced and thinning and loss of hair was also noted for high-dose males. Decrease in hemoglobin and erythrocytes at 0.4 and 2 mg/kg. SGOT, SGPT and alkaline phosphatase increased at 2 mg/kg.	incidence of subcutaneous masses and/or nodules; liver discoloration and nodules; atrophy of testes, seminal vesicles, and prostate. Increased incidence of mammary neoplasms (adenomas and adenocarcinomas).
Mice 50/sex/dose	0; 0.04; 0.4, and 2 mg/kg 105 weeks oral	No dose-related mortality. Thinning and loss of hair at 2 mg/kg. Slightly reduced body weight gain at 2 mg/kg.	Slightly increased incidence of skin masses and/or nodules and alopecia. No drug-related inflammatory, degenerative, proliferative and/or neoplastic lesions.
Dogs Beagle 4/sex/dose	0; 10; 32, and 100 mg/kg 55 weeks oral	No mortality. Excessive lacrimation, retarded pupillary reflex, mild conjunctivitis, hyperemia of gums, abdominal distention, sparsity of hair, and quieted behaviour. Laboratory tests: slightly elevated alkaline phosphatase and SGPT at 100 mg/kg in 2 dogs. Elevated sedimentation rate, slightly reduced lymphocytes with increase in segmented neutrophils and decrease in eosinophils.	Reduced adrenal, testes, and prostate weight for all cyproterone acetate-treated animals. Ovary and uterus weights reduced at 100 mg/kg. Liver weight slightly increased for some dogs. Histopathology: marked adrenal atrophy of zona fasciculata and reticularis, testicular atrophy and absence of spermatogenesis, some Leydig cell hyperplasia, prostatic atrophy, ovarian and uterine atrophy, hyperplasia of mammary gland in males and females.
Rhesus monkey 4 females/ dose	0; 0.04; 0.4, and 40 mg/kg 12 weeks oral	No mortality or behaviour changes. Dose-related alopecia. Raised insulin level above 0.04 mg/kg. Negative influence on	At doses of 0.4 mg/kg and above - diffuse liver cell hypertrophy and an increase in smooth endoplasmic reticulum.

Animal Species	Dosage and Duration	Mortality and Clinical and Laboratory Observations	Necropsy and Histopathology
		coagulation at 0.4 mg/kg and 40 mg/kg. Stimulation of ACTH cells at 0.4 mg and above. Increase in prolactin cells and slight reduction in gonadotrophin cells. Galactorrhea in all treated.	At the two highest doses, 2 and 3 animals also had occasional eosinophil cytoplasmic inclusion bodies in the liver cells. In most treated animals small mammary nodules were palpable in the glandular tissue; at 40 mg/kg slight ductus proliferation was also noted.

### **Mutagenesis and Carcinogenesis**

Recognized first-line tests of genotoxicity gave negative results when conducted with CPA. No mutagenic effect of cyproterone acetate was demonstrated in either *in vitro* (*Salmonella typhimurium*) or *in vivo* (micronucleus test in the monkey). However, further tests showed that CPA was capable of producing DNA adducts and an increase in DNA repair activity in liver cells from rats and monkeys and also in freshly isolated human hepatocytes.

This DNA-adduct formation occurred at systemic exposures that might be expected to occur in the recommended dose regimens for CPA. One *in vivo* consequence of CPA treatment was the increased incidence of focal, possible preneoplastic liver lesions in which cellular enzymes were altered in female rats. An initiating potential besides promoting effect of cyproterone acetate on the formation of ATPase deficient and GT positive foci in female rat livers was noted. CPA also enhanced the frequencies of mutations in the livers of female transgenic rats in a dose-dependent manner, indicating that CPA is mutagenic.

Investigations into the tumorigenicity of cyproterone acetate did not reveal a specific tumorigenic potential in the liver of rodents although other neoplasms, including mammary adenocarcinoma in rats, were observed (see Table 13).

### **Reproductive Toxicology**

Testicular atrophy and absence of spermatogenesis, some Leydig cell hyperplasia, prostatic atrophy, ovarian and uterine atrophy were observed in beagle dogs. A reduced number of pregnancies in untreated female rats was observed when male rats were administered with 40 mg/kg/day cyproterone acetate. The temporary inhibition of fertility in male rats brought about by daily oral treatment of cyproterone acetate did not result in malformations or impairment of fertility in the offspring produced by untreated female animals.

The treatment of pregnant animals with cyproterone acetate leads to developmental disturbances

in male fetuses. Testosterone-dependent differentiation processes are affected: signs of feminization of varying degrees of severity develop.

**Table 14: Fertility and Reproduction Study**

Animal Species	Route and Dosage of Administration	Findings
Rats 24/sex/dose (2 generations)	0; 0.4; 4.0 and 40 mg/kg oral	0.4 mg/kg: No influence by drug on fertility of the P1 and F1 generations. 4 mg/kg: Significant decrease in body weights but no impairment of pre- and postnatal development. 40 mg/kg: Food intake and body weight gain reduced. Although attempted matings were increased, less than 50% of the females had litters. No specific pathological changes were found in the dams, fetuses, or young. Similarly, no malformations were observed.

## REFERENCES

Apotex Research Inc. Comparative, Randomized, 2-Way Crossover Bioavailability Study of Cyproterone Acetate (Novex Pharma) and Androcur® (Berlex Canada Inc.) 50 mg Tablets/Under Fasting Conditions. Study Code CY0360. Data on File.

Bracci U, Di Silverio F. Present procedure with regard to the use of hormone therapy in cases of cancer of the prostate. In: Steffanini P, Speranza V, editors. Proc. 18, World Congr Int Coll Surg. Rome: Excerpta med., (Amst.) Int. Congr. Ser.; 1972. p. 275.

Bruchovsky N, Goldenberg SL, Coppin C, Brown E, McLoughlin M. Cyproterone acetate plus mini-dose diethylstilbestrol: initial hormone therapy for advanced prostatic cancer. In: Poster presentation at Satellite Symposium; 1984; Montreal; 1984.

Di Silverio F, Galiardi V. Il cancro della prostata. Nostri orientamenti terapeutici nelle forme estrogeno-resistenti. Boll. Atti Urol 1968; 5: 2.

Froelich S, Dali-Youcef N, Boyer P, Kehrl P, Maitrot D, Auwerx J, et al. Does cyproterone acetate promote multiple meningiomas? European Congress of Endocrinology 2008; Berlin, Germany: BioScientifica; 2008. p. 158.

Geller J, Vazakas G, Fruchtman B, Newman H, Nakao K, Loh A. The effect of cyproterone acetate on advanced carcinoma of the prostate. Surg Gynecol Obstet 1968;127(4):748-58.

Giuliani L, Pescatore D, Giberti C, Martorana G, Natta G. Treatment of advanced prostatic carcinoma with cyproterone acetate and orchiectomy--5-year follow-up. Eur Urol 1980;6(3):145-8.

Goldenberg SL, Bruchovsky N. Use of cyproterone acetate in prostate cancer. Urol Clin North Am 1991;18(1):111-22.

Goldenberg SL, Gleave ME. Current perspectives on the expanding role of Androcur®. In: Pharmanual: Pharma Libri Publishers Inc.; 1994.

Huan SD, Gerridzen RG, Yau JC, Stewart DJ. Antiandrogen withdrawal syndrome with nilutamide. Urology 1997;49(4):632-4.

Jacobi GH, Altwein JE, Kurth KH, Basting R, Hohenfellner R. Treatment of advanced prostatic cancer with parenteral cyproterone acetate: a phase III randomised trial. Br J Urol 1980;52(3):208-15.

Laufer M, Sinibaldi VJ, Carducci MA, Eisenberger MA. Rapid disease progression after the administration of bicalutamide in patients with metastatic prostate cancer. *Urology* 1999;54(4):745.

Longmore L, Foley JP, Rozanski TA, Higgins B, Thompson IM. Prolonged prostate-specific antigen response in flutamide withdrawal syndrome despite disease progression. *South Med J* 1998;91(6):573-5.

Mellin P. Cyproterone therapy in prostatic cancer stage IV. In: Raspé G, Brosig W, editors. *International Symposium on the Treatment of Carcinoma of the Prostate (1969 Berlin, Germany)*. Life sciences monographs. Vieweg, Oxford, Edinburgh, New York, Toronto, Sydney, Brauchweig: Pergamon Press; 1971. p. 180.

Morse HC, Leach DR, Rowley MJ, Heller CG. Effect of cyproterone acetate on sperm concentration, seminal fluid volume, testicular cytology and levels of plasma and urinary ICSH, FSH and testosterone in normal men. *J Reprod Fertil* 1973;32(3):365-78.

Nagamatsu GR. Use of cyproterone acetate in stage III and IV. In: Raspé G, Brosig W, editors. *International Symposium on the Treatment of Carcinoma of the Prostate (1969 Berlin, Germany)*. Life sciences monographs. Vieweg, Oxford, Edinburgh, New York, Toronto, Sydney, Brauchweig: Pergamon Press; 1971. p. 173.

Neumann F. The antiandrogen cyproterone acetate: discovery, chemistry, basic pharmacology, clinical use and tool in basic research. *Exp Clin Endocrinol* 1994;102(1):1-32. Nieuweboer B, Lubke K. Radioimmunological determination of cyproterone acetate. *Horm Res* 1977;8(4):210-8.

Pavone M, et al. Medroxyprogesterone acetate, Diethylstilboestrol and Cyproterone Acetate in the treatment of prostatic cancer. Interim report of a prospective study of the European Organization for Research on the Treatment of Cancer (EORTC) Genito-Urinary Tract Cooperative group. *Excerpt. Med. Int. Congr. Ser.* 1982;677:436-44.

Peetermans C, Roux E. Androcur 50 in the treatment of prostatic carcinoma. Belgian multicentric study with the participation of 30 urologists. *Acta Urol Belg* 1994;62(1):77-82.

Pescatore D, Giberti C, Martorana G, Natta G, Giuliani L. The effect of cyproterone acetate and orchiectomy on metastases from prostatic cancer. *Eur Urol* 1980;6(3):149-53.

Prostate Cancer Trialists' Collaborative Group. Maximum androgen blockade in advanced prostate cancer: an overview of the randomised trials. *Prostate Cancer Trialists' Collaborative Group. Lancet.* 2000;355(9214):1491-8.

Schuppler J, Gunzel P. Cyproterone acetate. *Lancet* 1976;7987:688.

Scott WW, Schirmer HK. A new oral progestational steroid effective in treating prostatic cancer. *Trans Am Assoc Genitourin Surg* 1966;58:54-62.

Scott WW, Wade JC. Medical treatment of benign nodular prostatic hyperplasia with cyproterone acetate. *J Urol* 1969;101(1):81-5.

Sella A, Flex D, Sulkes A, Baniel J. Antiandrogen withdrawal syndrome with cyproterone acetate. *Urology* 1998;52(6):1091-3.

Speck U, Jentsch D, Kuhne G, Schulze PE, Wendt H. [Bioavailability and pharmacokinetics of <sup>14</sup>C-cyproterone acetate after administration as a 50-mg tablet (author's transl)]. *Arzneimittelforschung* 1976;26(9):1717-20.

Varenhorst E, Risberg B. Effects of estrogen, orchidectomy, and cyproterone acetate on tissue fibrinolysis in patients with carcinoma of the prostate. *Invest Urol* 1981;18(5):355-7.

Wirth MP, Froschermaier SE. The antiandrogen withdrawal syndrome. *Urol Res* 1997;25 Suppl 2:S67-71.

Product Monograph - ANDROCUR<sup>®</sup> (cyproterone acetate tablets) 50 mg  
ANDROCUR<sup>®</sup> DEPOT (cyproterone acetate injection) 100 mg/mL, Bayer Inc., Date of  
Revision: February 25, 2011. Control Number: 142634

**PART III: CONSUMER INFORMATION****PrCYPROTERONE**

Cyproterone Acetate Tablets  
House Standard

*This leaflet is Part 3 of a three-part "Product Monograph" published when CYPROTERONE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about CYPROTERONE. Contact your doctor or pharmacist if you have any questions*

**ABOUT THE DRUG. ABOUT THIS MEDICATION****What the medication is used for:**

CYPROTERONE is indicated for the palliative treatment of patients with advanced prostate cancer.

**What it does:**

CYPROTERONE is a drug acting against male sex hormones (androgens). Androgens encourage the growth of prostate cancer and CYPROTERONE inhibits this effect.

**When it should not be used:**

Do not take CYPROTERONE if you:

- are allergic (hypersensitive) to cyproterone acetate or any of the other ingredients of CYPROTERONE;
- have a liver disease or reduced liver function;
- have Dubin-Johnson syndrome or Rotor syndrome that affects the excretion of red blood cell pigment (bilirubin);
- have or have had liver tumors that are not due to the spread of prostate cancer;
- have or ever had a benign brain tumor (meningioma)
- have wasting diseases that are not related to prostate cancer;
- suffer from severe chronic depression;
- have conditions that increase your risk for developing blood clots (thromboembolic process).

**What the medicinal ingredient is:**

Cyproterone acetate

**What the nonmedicinal ingredients are:**

The other tablet ingredients are colloidal silicon dioxide, croscarmellose sodium and magnesium stearate.

**What dosage forms it comes in:**

CYPROTERONE is supplied in a form of :  
Tablet: Each tablet contains 50 mg cyproterone acetate

**WARNINGS AND PRECAUTIONS****Serious Warnings and Precautions**

**CYPROTERONE should be prescribed and managed by a doctor experienced with the treatment of prostate cancer.**

**Possible serious side effects of CYPROTERONE include:**

- **Liver damage and liver failure**

BEFORE you use CYPROTERONE talk to your doctor or pharmacist if you have or have had any of the following conditions:

- Blood clots;
- Liver problem;
- Depression;
- Breathing problem (shortness of breath).
- Diabetes. Your doctor may need to adjust your antidiabetic medication. This requires strict supervision during treatment with CYPROTERONE.
- Anemia. Your doctor will monitor your red-blood cell count during treatment.
- History of benign brain tumors (meningiomas)

CYPROTERONE may decrease the function of the hormone-producing glands attached to the kidney (adrenal glands).

Benign brain tumours (single and multiple meningiomas) have been reported with long-term treatment of cyproterone acetate and very rarely in patients after several months of treatment with cyproterone acetate for prostate cancer.

CYPROTERONE is not recommended for use in children and adolescents below 18 years of age and before the end of puberty, as it may have a negative effect on growth and the immature hormonal regulation functions.

CYPROTERONE is not recommended for use in women.

Based on an analysis of the results of several older studies, long-term combination therapy of cyproterone acetate with either orchiectomy (removal of testicles) or a GnRH agonist (one class of drug that acts against male sex hormones) in advanced prostate cancer may result in a 5-year survival disadvantage compared to castration alone.

## INTERACTIONS WITH THIS MEDICATION

Drugs that may interact with CYPROTERONE include:

- Statins (medicines for reducing blood fats)
- Ketoconazole, itraconazole, clotrimazole (for fungal infections)
- Ritonavir (for viral infections)
- Rifampicin (for tuberculosis)
- Phenytoin (for epilepsy)
- St. John's Wort (herbal remedy for depression)

## PROPER USE OF THIS MEDICATION

**You should follow the dose prescribed by your doctor.**

### Usual dose

#### **CYPROTERONE Tablet**

200 mg to 300 mg daily taken orally in two or three divided doses with liquid after meals. Maximum daily dose is 300 mg. After orchiectomy (removal of testicles), a lower daily dose of 2 to 4 tablets (100-200 mg) is recommended.

### **Overdose:**

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

### **Missed Dose**

#### **CYPROTERONE Tablet**

Do not take the missed CYPROTERONE tablet (do not take a double dose to make up for the missed tablet). Continue taking the tablets at the regular time.

## SIDE EFFECTS AND WHAT TO DO ABOUT THEM

CYPROTERONE can have side effects, like all medicines. For further information about any of these effects, ask a doctor or pharmacist.

If any of the side effects get serious, or if you notice any side effects not listed here, please tell your doctor or pharmacist.

Very frequent side effects:

- Reduced sexual interest (decreased libido)
- Swelling of the breast, breast soreness (gynecomastia)
- Impotence
- Abnormal sperm, low sperm count which may be reversible after CYPROTERONE is discontinued.

Other side effects:

- Hair loss or unusual increase in hair growth
- Constipation or diarrhea (loose stools)
- Weight gain
- Tiredness and weakness
- Shortness of breath

- Skin rash, blisters
- Vision change
- Decrease in blood pressure
- Increase in blood sugar
- Depression
- Pulmonary oil microembolism with symptoms such as cough, shortness of breath (dyspnea), or chest pain
- General postinjection reactions (vasovagal reactions) with symptoms such as malaise, increased sweating (hyperhidrosis), dizziness, pins and needles (parasthesia) or fainting (syncope)

Rare but serious side effects:

- Liver toxicity: generally feeling unwell, fever, nausea, vomiting, loss of appetite, itching, yellowing of the skin and eyes, light-colored stools, dark-colored urine
- Life-threatening internal bleeding (intra-abdominal hemorrhage): unusual upper abdominal pains which do not disappear within a short time
- Blood clots: swelling of the calf or leg (blood clots in the leg), chest pain and being short of breath (blood clots in the lung), suddenly feeling weak, loss of coordination, slurred speech (a stroke or blood clots in the brain), temporary blindness (blood clots in the eye)

Bone loss (osteoporosis) and benign brain tumors (cerebral meningioma) have been reported with long term use of cyproterone acetate .

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptom / effect	Talk with your doctor		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Inability to achieve or maintain an erection	☺		
Liver toxicity, (generally feeling unwell, fever, nausea, vomiting, loss of appetite, itching all over the body, yellowing of the skin or eyes, light colored bowel movements, dark urine) liver inflammation (hepatitis), liver failure			☺
Reduced sexual drive	☺		
Reversible inhibition of sperm production	☺		
Swelling of the calf or leg, chest pain, suddenly feeling weak			☺

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor	Stop taking drug and
Unusual upper abdominal pains which do not disappear within a short time		ç

This is not a complete list of side effects. For any unexpected effects while taking CYPROTERONE, contact your doctor or pharmacist.

**HOW TO STORE IT**

Store between 15 and 30°C. Keep container tightly closed, Keep out of the reach and sight of children.

Do not take CYPROTERONE after the expiry date which is stated on the pack.

Medicines must not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicine no longer required. These measures will help to protect the environment.

**Reporting Suspected Side Effects**

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Fax toll-free to 1-866-678-6789, or
  - Mail to:  
Canada Vigilance Program  
Health Canada  
Postal Locator 0701E  
Ottawa, Ontario  
K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

*NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

For more information, please contact your doctor, pharmacist or other healthcare professional.

This leaflet plus the full product monograph, prepared for health professionals, can be obtained by contacting the sponsor, AA Pharma Inc. at:

1-877-998-9097

This leaflet was prepared by AA Pharama Inc., Vaughan, Ontario, L4K 4N7

Last revised: November 25, 2014